

Peer-Reviewed Publications Using Pinnacle Technology's Products
(exception is paper by Hu et al that provides an overview of the sensor technology)

Biosensors

- Agnesi, F., Blaha, C.D., Lin, J., & Lee, K.H. (2010). Local glutamate release in the rat ventral lateral thalamus evoked by high-frequency stimulation. *Journal of Neural Engineering*, 7(2). [doi:10.1088/1741-2560/7/2/026009](https://doi.org/10.1088/1741-2560/7/2/026009)
- Agnesi, F., Tye, S.J., Bledsoe, J.M., Griessenauer, C.J., Kimble, C.J., Sieck, G.C., Bennet, K.E., Garris, P.A., Blaha, C.D., & Lee, K.H. (2009, May 8). Wireless instantaneous neurotransmitter concentration system-based amperometric detection of dopamine, adenosine, and glutamate for intraoperative neurochemical monitoring. *Journal of Neurosurgery*. [doi:10.3171/2009.3.JNS0990](https://doi.org/10.3171/2009.3.JNS0990)
- Behrend, C.E., Cassim, S.M., Pallone, M.J., Daubenspeck, J.A., Hartov, A., Roberts, D.W., & Leiter, J.C. (2009). Toward feedback controlled deep brain stimulation: Dynamics of glutamate release in the subthalamic nucleus in rats. *Journal of Neuroscience Methods*, 180(2): 278-289. [doi:10.1016/j.jneumeth.2009.04.001](https://doi.org/10.1016/j.jneumeth.2009.04.001)
- Bonaventure, P., Aluisio, L., Shoblock, J., Boggs, J.D., Fraser, I.C., Lord, B., Lovenberg, T.W., & Galici, R. (2011). Pharmacological blockade of serotonin 5-HT₇ receptor reverses working memory deficits in rats by normalizing cortical glutamate neurotransmission. *PLoS ONE*, 6(6): 1-7. [doi:10.1371/journal.pone.0020210](https://doi.org/10.1371/journal.pone.0020210)
- Cairns, B.E., Dong, X., Mann, M.K., Svensson, P., Sessle, B.J., Arendt-Nielsen, L., & McErlane, K.M. (2007). Systemic administration of monosodium glutamate elevates intramuscular glutamate levels and sensitizes rat masseter muscle afferent fibers. *Pain*, 132: 33-41. [doi:10.1016/j.pain.2007.01.023](https://doi.org/10.1016/j.pain.2007.01.023)
- Chang, S., Shon, Y.M., Agnesi, F., & Lee, K.H. (2009, September). Microthalamotomy effect during deep brain stimulation: Potential involvement of adenosine and glutamate efflux. Engineering in Medicine and Biology Society. Annual International Conference of the IEEE, 3294-3297. [doi:10.1109/IEMBS.2009.5333735](https://doi.org/10.1109/IEMBS.2009.5333735)
- Endo, H., Takahashi, E., Murata, M., Ohnuki, H., Ren, H., Tsugawa, W., & Sode, K. (2010). Wireless monitoring of blood glucose levels in flatfish with a needle biosensor. *Fisheries Science*, 76(4): 687-694. [doi:10.1007/s12562-010-0256-0](https://doi.org/10.1007/s12562-010-0256-0)
- Endo, H., Yonemori, Y., Hibi, K., Ren, H., Hayashi, T., Tsugawa, W., & Sode, K. (2009). Wireless enzyme sensor system for real-time monitoring of blood glucose levels in fish. *Biosensors Bioelectronics*, 24(5): 1417-1423. [doi:10.1016/j.bios.2008.08.038](https://doi.org/10.1016/j.bios.2008.08.038)
- Gass, J.T., Sinclair, C.M., Cleva, R.M., Widholm, J.J., & Olive, M.F. (2011). Alcohol-seeking behavior is associated with increased glutamate transmission in basolateral amygdala and nucleus accumbens as measured by glutamate-oxidase-coated biosensors. *Addiction Biology*, 16(2): 215-228. [doi:10.1111/j.1369-1600.2010.00262.x](https://doi.org/10.1111/j.1369-1600.2010.00262.x)
- Gazerani, P., Au, S., Dong, X., Kumar, U., Arendt-Nielsen, L., & Cairns, B.E. (2010). Botulinum neurotoxin type A (BoNTA) decreases the mechanical sensitivity of nociceptors and inhibits neurogenic vasodilation in a craniofacial muscle targeted for migraine prophylaxis. *Pain*, 151(3): 606-616. [doi:10.1016/j.pain.2010.07.029](https://doi.org/10.1016/j.pain.2010.07.029)
- Gifford, R., Kehoe, J.J., Barnes, S.L., Kornilayev, B.A., Alterman, M.A., & Wilson, G.S. (2006). Protein interactions with subcutaneously implanted biosensors. *Biomaterials*, 27(12): 2587-2598. [doi:10.1016/j.biomaterials.2005.11.033](https://doi.org/10.1016/j.biomaterials.2005.11.033)
- Gifford, R., Batchelor, M.M., Lee, Y., Gokulrangan, G., Meyerhoff, M.E., & Wilson, G.S. (2005). Mediation of *in vivo* glucose sensor inflammatory response via nitric oxide release. *Journal of Biomedical Materials Research Part A*, 75(4): 755-766. [doi:10.1002/jbm.a.30359](https://doi.org/10.1002/jbm.a.30359)
- Hakim, A.W., Dong, X., & Cairns, B.E. (2011). TNF α mechanically sensitizes masseter muscle nociceptors by increasing prostaglandin E₂ levels. *Journal of Neurophysiology*, 105(1): 154-161. [doi:10.1152/jn.00730.2010](https://doi.org/10.1152/jn.00730.2010)
- Hu, Y., Mitchell, K.M., Albahadily, F.N., Michaelis, E.K., & Wilson, G.S. (1994). Direct measurement of glutamate release in the brain using a dual enzyme-based electrochemical sensor. *Brain Research*, 659(1-2): 117-125. [doi:10.1016/0006-8993\(94\)90870-2](https://doi.org/10.1016/0006-8993(94)90870-2)
- John, J., Ramanathan, L., & Siegel, J.M. (2008). Rapid changes in glutamate levels in the posterior hypothalamus across sleep-wake states in freely behaving rats. *American Journal of Physiology: Regulatory, Integrative, and Comparative Physiology*, 295(6): R2041-R2049. [doi:10.1152/ajpregu.90541.2008](https://doi.org/10.1152/ajpregu.90541.2008)

- Lee, K.H., Hitti, F.L., Chang, S., Lee, D.C., Roberts, D.W., McIntyre, C.C., & Leiter, J.C. (2011). High frequency stimulation abolishes thalamic network oscillations: An electrophysiological and computational analysis. *Journal of Neural Engineering*, 8(4): 1-11. [doi:10.1088/1741-2560/8/4/046001](https://doi.org/10.1088/1741-2560/8/4/046001)
- Lee, K.H., Kristic, K., van Hoff, R., Hitti, F.L., Blaha, C., Harris, B., Roberts, D.W., & Leiter, J.C. (2007). High-frequency stimulation of the subthalamic nucleus increases glutamate in the subthalamic nucleus of rats as demonstrated by in vivo enzyme-linked glutamate sensor. *Brain Research*, 1162(8): 121-129. [doi:10.1016/j.brainres.2007.06.021](https://doi.org/10.1016/j.brainres.2007.06.021)
- Matsuno, A., & Inoue, H. (2008). Hippocampal glutamate release on learning and memory in teeth-loss rats. *Prosthodontic Research & Practice*, 7(2): 71-77. [doi:10.2186/prp.7.71](https://doi.org/10.2186/prp.7.71)
- Morita, H., Abe, C., Awazu, C., & Tanaka, K. (2007). Long-term hypergravity induces plastic alterations in vestibulo-cardiovascular reflex in conscious rats. *Neuroscience Letters*, 412(3): 201-205. [doi:10.1016/j.neulet.2006.11.014](https://doi.org/10.1016/j.neulet.2006.11.014)
- Morrell, C.N., Sun, H., Ikeda, M., Beique, J., Swaim, A.M., Mason, E., Martin, T.V., Thompson, L.E., Gozen, O., Ampagoomian, D., Sprengel, R., Rothstein, J., Faraday, N., Huganir, R., & Lowenstein, C.J. (2008). Glutamate mediates platelet activation through the AMPA receptor. *Journal of Experimental Medicine*, 205(3): 575-584. [doi:10.1084/jem.20071474](https://doi.org/10.1084/jem.20071474)
- Otsuguro, K., Wada, M., & Ito, S. (2011). Differential contributions of adenosine to hypoxia-evoked depressions of three neuronal pathways in isolated spinal cord of neonatal rats. *British Journal of Pharmacology*, 164(1): 132-144. [doi:10.1111/j.1476-5381.2011.01333.x](https://doi.org/10.1111/j.1476-5381.2011.01333.x)
- Takaoka, H., & Yasuzawa, M. (2010). Fabrication of an implantable fine needle-type glucose sensor using γ -polyglutamic acid. *Analytical Sciences*, 26(5): 551-555. [doi:10.2116/analsci.26.551](https://doi.org/10.2116/analsci.26.551)
- Tawfik, V.L., Chang, S., Hitti, F.L., Roberts, D.W., Leiter, J.C., Jovanovic, S., & Lee, K.H. (2010). Deep brain stimulation results in local glutamate and adenosine release: Investigation into the role of astrocytes. *Neurosurgery*, 67(2): 367-375. [doi:10.1227/01.NEU.0000371988.73620.4C](https://doi.org/10.1227/01.NEU.0000371988.73620.4C)
- Uslaner, J.M., Smith, S.M., Huszar, S.L., Pachmerhiwala, R., Hinchliffe, R.M., Vardigan, J.D., Nguyen, S.J., Surles, N.O., Yao, L., Barrow, J.C., Uebele, V.N., Renger, J.J., Clark, J., & Hutson, P.H. (2010). T-type calcium channel antagonism produces antipsychotic-like effects and reduces stimulant-induced glutamate release in the nucleus accumbens of rats. *Neuropharmacology*. Advance online publication. [doi:10.1016/j.neuropharm.2010.11.015](https://doi.org/10.1016/j.neuropharm.2010.11.015)
- Van Gompel, J.J., Chang, S., Goerss, S.J., Kim, I.Y., Kimble, C., Bennet, K.E., & Lee, K.H. (2010). Development of intraoperative electrochemical detection: Wireless instantaneous neurochemical concentration sensor for deep brain stimulation feedback. *Neurosurgical FOCUS*, 29(2): E6. [doi:10.3171/2010.5.FOCUS10110](https://doi.org/10.3171/2010.5.FOCUS10110)
- Yonemori, Y., Takahashi, E., Ren, H., Hayashi, T., & Endo, H. (2009). Biosensor system for continuous glucose monitoring in fish. *Analytica Chimica Acta*, 633(1): 90-96. [doi:10.1016/j.aca.2008.11.023](https://doi.org/10.1016/j.aca.2008.11.023)
- Yoneyama, Y., Yonemori, Y., Murata, M., Ohnuki, H., Hibi, K., Hayashi, T., Ren, H., & Endo, H. (2009). Wireless biosensor system for real-time cholesterol monitoring in fish "Nile tilapia." *Talanta*, 80(2): 909-915. [doi:10.1016/j.talanta.2009.08.014](https://doi.org/10.1016/j.talanta.2009.08.014)

Sleep & Biosensor

- Naylor, E., Aillon, D.V., Gabbert, S., Harmon, H., Johnson, D.A., Wilson, G.S., & Petillo, P.A. (2011). Simultaneous real-time measurement of EEG/EMG and L-glutamate in mice: A biosensor study of neuronal activity during sleep. *Journal of Electroanalytical Chemistry*, 656(1-2): 106-113. [doi:10.1016/j.jelechem.2010.12.031](https://doi.org/10.1016/j.jelechem.2010.12.031)

Sleep

- Altimus, C.M., Guler, A.D., Villa, K.L., McNeill, D.S., LeGates, T.A., & Hattar, S. (2008). Rods-cones and melanopsin detect light and dark to modulate sleep independent of image formation. *Proceedings of the National Academy of Sciences USA*, 105(50): 19998-20003. [doi:10.1073/pnas.0808312105](https://doi.org/10.1073/pnas.0808312105)
- Anaclet, C., Pedersen, N.P., Fuller, P.M., & Lu, J. (2010). Brainstem circuitry regulating phasic activation of trigeminal motoneurons during REM sleep. *PLoS ONE*, 5(1): e8788. [doi:10.1371/journal.pone.0008788](https://doi.org/10.1371/journal.pone.0008788)

October 24, 2011

- Gerashchenko, D., Wisor, J.P., Burns, D., Reh, R.K., Shiromani, P.J., Sakurai, T., de la Iglesia, H.O., & Kilduff, T.S. (2008). Identification of a population of sleep-active cerebral cortex neurons. *Proceedings of the National Academy of Sciences USA*, 105(29): 10227-10232. doi: [10.1073/pnas.0803125105](https://doi.org/10.1073/pnas.0803125105)
- Krenzer, M., Anaclet, C., Vetrivelan, R., Wang, N., Vong, L., Lowell, B.B., Fuller, P.M., Lu, J. (2011). Brainstem and spinal cord circuitry regulating REM sleep and muscle atonia. *PLoS ONE*, 6(10): e24998. doi: [10.1371/journal.pone.0024998](https://doi.org/10.1371/journal.pone.0024998)
- Narita, M., Niikura, K., Nanjo-Niikura, K., Narita, M., Furuya, M., Yamashita, A., Saeki, M., Matsushima, Y., Imai, S., Shimizu, T., Asato, M., Kuzumaki, N., Okutsu, D., Miyoshi, K., Suzuki, M., Tsukiyama, Y., Konno, M., Yomiya, K., Matoba, M., & Suzuki, T. (2011). Sleep disturbances in a neuropathic pain-like condition in the mouse are associated with altered GABAergic transmission in the cingulate cortex. *Pain*, 152(6): 1358-1372. doi: [10.1016/j.pain.2011.02.016](https://doi.org/10.1016/j.pain.2011.02.016)
- Rahmadi, M., Narita, M., Yamashita, A., Imai, S., Kuzumaki, N., & Suzuki, T. (2011). Sleep disturbance associated with an enhanced orexinergic system induced by chronic treatment with paroxetine and milnacipran. *Synapse*, 65(7): 652-657. doi: [10.1002/syn.20893](https://doi.org/10.1002/syn.20893)
- Sheward, W.J., Naylor, E., Knowles-Barley, S., Armstrong, J.D., Brooker, G.A., Seckl, J.R., Turek, F.W., Holmes, M.C., Zee, P.C., & Hargmar, A.J. (2010). Circadian control of mouse heart rate and blood pressure by the suprachiasmatic nuclei: Behavioral effects are more significant than direct outputs. *PLoS ONE*, 5(3): e9783. doi: [10.1371/journal.pone.0009783](https://doi.org/10.1371/journal.pone.0009783)
- Takemura, Y., Yamashita, A., Horiuchi, H., Furuya, M., Yanase, M., Niikura, K., Imai, S., Hatakeyama, N., Kinoshita, H., Tsukiyama, Y., Senba, E., Matoba, M., Kuzumaki, N., Yamazaki, M., Suzuki, T., & Narita, M. (2011). Effects of gabapentin on brain hyperactivity related to pain and sleep disturbance under a neuropathic pain-like state using fMRI and brain wave analysis. *Synapse*, 65(7): 668-676. doi: [10.1002/syn.20898](https://doi.org/10.1002/syn.20898)
- Seizure**
- Baraban, S., Southwell, D.G., Estrada, R.C., Jones, D.L., Sebe, J.Y., Alfaro-Cervello, C., Garcia-Verdugo, J.M., Rubenstein, J.L.R., & Alvarez-Buylla, A. (2009). Reduction of seizures by transplantation of cortical GABAergic interneuron precursors into Kv1.1 mutant mice. *Proceedings of the National Academy of Sciences USA*, 106(36): 15472-15477. doi: [10.1073/pnas.0900141106](https://doi.org/10.1073/pnas.0900141106)
- Bell, M.E., & Bialecki, R. (2009). Differences in electrographic ICTAL activity between C57BL/6 and BALB/c mouse strains. *Journal of Pharmacological Toxicological Methods*, 60(2):229. doi: [10.1016/j.vascn.2009.04.089](https://doi.org/10.1016/j.vascn.2009.04.089)
- Calderon, D.P., Fremont, R., Kraenzlin, F., & Khodakhah, K. (2011). The neural substrates of rapid-onset Dystonia-Parkinsonism. *Nature Neuroscience*, 14: 357-365. doi: [10.1038/nn.2753](https://doi.org/10.1038/nn.2753)
- Chung, W.K., Shin, M., Jaramillo, T.C., Leibel, R.L., LeDuc, C.A., Fischer, S.G., Tzilianos, E., Gheith, A.A., Lewis, A.S., & Chetkovich, D.M. (2009). Absence epilepsy in *apathetic*, a spontaneous mutant mouse lacking the h channel subunit, HCN2. *Neurobiology of Disease*, 33(3): 499-508. doi: [10.1016/j.nbd.2008.12.004](https://doi.org/10.1016/j.nbd.2008.12.004)
- Cuaycong, M., Engel, M., Weinstein, S.L., Salmon, E., Perlman, J.M., Sunderam, S., & Vannucci, S.J. (2011). A novel approach to the study of hypoxia-ischemia-induced clinical and subclinical seizures in the neonatal rat. *Developmental Neuroscience*. doi: [10.1159/000331646](https://doi.org/10.1159/000331646)
- Edwards, D.A., Shah, H.P., Cao, W., Gravenstein, N., Suebert, C.N., & Martynyuk, A.E. (2010). Bumetanide alleviates epileptogenic and neurotoxic effects of sevoflurane in neonatal rat brain. *Anesthesiology*, 112(3): 567-575. doi: [10.1097/ALN.0b013e3181cf9138](https://doi.org/10.1097/ALN.0b013e3181cf9138)
- Galanopoulou, A.S. (2008). Dissociated gender-specific effects of recurrent seizures on GABA signaling in CA1 pyramidal neurons: Role of GABA_A receptors. *Journal of Neuroscience*, 28(7): 1557-1567. doi: [10.1523/JNEUROSCI.5180-07.2008](https://doi.org/10.1523/JNEUROSCI.5180-07.2008)
- Greenwood, J.S.F., Wang, Y., Estrada, R.C., Ackerman, L., Ohara, P.T., & Baraban, S.C. (2009). Seizures, enhanced excitation, and increased vesicle number in *Lis1* mutant mice. *Annals of Neurology*, 66(5): 644-653. doi: [10.1002/ana.21775](https://doi.org/10.1002/ana.21775)
- Hawkins, N.A., Martin, M.S., Frankel, W.N., Kearney, J.A., & Escayg, A. (2011). Neuronal voltage-gated ion channels are genetic modifiers of generalized epilepsy with febrile seizures plus. *Neurobiology of Disease*, 41(3): 655-660. doi: [10.1016/j.nbd.2010.11.016](https://doi.org/10.1016/j.nbd.2010.11.016)
- Hitt, B.D., Jaramillo, T.C., Chetkovich, D.M., & Vassar, R. (2010). BACE1^{-/-} mice exhibit seizure activity that does not correlate with sodium channel level or axonal localization. *Molecular Neurodegeneration*, 5(31). doi: [10.1186/1750-1326-5-31](https://doi.org/10.1186/1750-1326-5-31)

- Hu, X., Zhou, X., He, W., Yang, J., Xiong, W., Wong, P., Wilson, C.G., & Yan, R. (2010). BACE1 deficiency causes altered neuronal activity and neurodegeneration. *Journal of Neuroscience*, 30(26): 8819-8829. [doi:10.1523/JNEUROSCI.1334-10.2010](https://doi.org/10.1523/JNEUROSCI.1334-10.2010)
- Lehnart, S.E., Mongillo, M., Bellinger, A., Lindegger, N., Chen, B., Hsueh, W., Reiken, S., Wronska, A., Drew, L.J., Ward, C.W., Lederer, W.J., Kass, R.S., Morley, G., & Marks, A.R. (2008). Leaky Ca²⁺ release channel/ryanodine receptor 2 causes seizures and sudden cardiac death in mice. *Journal of Clinical Investigation*, 118(6): 2230-2245. [doi:10.1172/JCI35346](https://doi.org/10.1172/JCI35346)
- Marchi, N., Fan, Q., Ghosh, C., Fazio, V., Bertolini, F., Betto, G., Batra, A., Carlton, E., Najm, I., Granata, T., & Janigro, D. (2009). Antagonism of peripheral inflammation reduces the severity of status epilepticus. *Neurobiology of Disease*, 33(2): 171-181. [doi:10.1016/j.nbd.2008.10.002](https://doi.org/10.1016/j.nbd.2008.10.002)
- Marchi, N., Teng, Q., Ghosh, C., Fan, Q., Nguyen, M.T., Desai, N.K., Bawa, H., Rasmussen, P., Masaryk, T.K., & Janigro, D. (2010). Blood-brain barrier damage, but not parenchymal white blood cells, is a hallmark of seizure activity. *Brain Research*, 1353: 176-186. [doi:10.1016/j.brainres.2010.06.051](https://doi.org/10.1016/j.brainres.2010.06.051)
- Ono, T., Moshé, S.L., & Galanopoulou, A.S. (2011). Carisbamate acutely suppresses spasms in a rat model of symptomatic infantile spasms. *Epilepsia*. Advance online publication. [doi:10.1111/j.1528-1167.2011.03173.x](https://doi.org/10.1111/j.1528-1167.2011.03173.x)
- Raffo, E., Coppola, A., Ono, T., Briggs, S.W., & Galanopoulou, A.S. (2011). A pulse rapamycin therapy for infantile spasms and associated cognitive decline. *Neurobiology of Disease*, 43(2): 322-329. [doi:10.1016/j.nbd.2011.03.021](https://doi.org/10.1016/j.nbd.2011.03.021)
- Rajdev, P., Ward, M., & Irazoqui, P. (2011). Effect of stimulus parameters in the treatment of seizures by electrical stimulation in the kainate animal model. *International Journal of Neural Systems*, 21(2): 151-162. [doi:10.1142/S0129065711002730](https://doi.org/10.1142/S0129065711002730)
- Scantlebury, M.H., Galanopoulou, A.S., Chudomelova, L., Raffo, E., Betancourth, D., & Moshe, S.L. (2010). A model of symptomatic infantile spasms syndrome. *Neurobiology of Disease*, 37(3): 604-612. [doi:10.1016/j.nbd.2009.11.011](https://doi.org/10.1016/j.nbd.2009.11.011)
- Sebe, J.Y., & Baraban, S.C. (2010). The promise of an interneuron-based cell therapy for epilepsy. *Developmental Neurobiology*, 71(1): 107-117. [doi:10.1002/dneu.20813](https://doi.org/10.1002/dneu.20813)
- Shin, M., Brager, D., Jaramillo, T.C., Johnston, D., & Chetkovich, D.M. (2008). Mislocalization of h channel subunits underlies h channelopathy in temporal lobe epilepsy. *Neurobiology of Disease*, 32(1): 26-36. [doi:10.1016/j.nbd.2008.06.013](https://doi.org/10.1016/j.nbd.2008.06.013)
- Vogt, D.L., Thomas, D., Galvan, V., Bredesen, D.E., Lamb, B.T., & Pimplikar, S.W. (2011). Abnormal neuronal networks and seizure susceptibility in mice overexpressing the APP intracellular domain. *Neurobiology of Aging*, 32(9): 1725-1729. [doi:10.1016/j.neurobiolaging.2009.09.002](https://doi.org/10.1016/j.neurobiolaging.2009.09.002)
- Velišek, L., Shang, E., Velišková, J., Chachua, T., Macchiarulo, S., Maglakelidze, G., Wolgemuth, D.J., & Greenberg, D. (2011). GABAergic neuron deficit as an idiopathic generalized epilepsy mechanism: The role of BRD2 haploinsufficiency in juvenile myoclonic epilepsy. *PLoS ONE*, 6(8): e23656. [doi:10.1371/journal.pone.0023656](https://doi.org/10.1371/journal.pone.0023656)
- Wang, Y., Dye, C.A., Sohal, V., Long, J.E., Estrada, R.C., Roztocil, T., Lufkin, T., Deisseroth, K., Baraban, S.C., & Rubenstein, J.L.R. (2010). *Dlx5* and *Dlx6* regulate the development of parvalbumin-expressing cortical interneurons. *Journal of Neuroscience*, 30(15): 5334-5345. [doi:10.1523/JNEUROSCI.5963-09.2010](https://doi.org/10.1523/JNEUROSCI.5963-09.2010)

Other

- Fischer, D.J., Hulvey, M.K., Regel, A.R., & Lunte, S.M. (2009). Amperometric detection in microchip electrophoresis devices: Effect of electrode material and alignment on analytical performance. *Electrophoresis*, 30(19): 3324-3333. [doi:10.1002/elps.200900317](https://doi.org/10.1002/elps.200900317)
- Guisseppi-Elie, A. (2011). An implantable biochip to influence patient outcomes following trauma-induced hemorrhage. *Analytical and Bioanalytical Chemistry*, 399(1): 403-419. [doi:10.1007/s00216-010-4271-x](https://doi.org/10.1007/s00216-010-4271-x)
- Gunasekara, D.B., Hulvey, M.K., & Lunte, S.M. (2011). In-channel amperometric detection for microchip electrophoresis using a wireless isolated potentiostat. *Electrophoresis*, 32(8): 832-837. [doi:10.1002/elps/201000681](https://doi.org/10.1002/elps/201000681)
- Hodges, M.R., Tattersall, G.J., Harris, M.B., McEvoy, S.D., Richerson, D.N., Deneris, E.S., Johnson, R.L., Chen, Z.F., & Richerson, G.B. (2008). Defects in breathing and thermoregulation in mice with near-complete absence of central serotonin neurons. *Journal of Neuroscience*, 28(10): 2495-2505. [doi:10.1523/JNEUROSCI.4729-07.2008](https://doi.org/10.1523/JNEUROSCI.4729-07.2008)

Publications Using Pinnacle's Products

October 24, 2011

- Kiyatkin, E.A., & Smirnov, M.S. (2009). Rapid EEG desynchronization and EMG activation induced by intravenous cocaine in freely moving rats: a peripheral, nondopamine neural triggering. *American Journal of Physiology: Regulatory, Integrative, and Comparative Physiology*, 298(2): R285-R300. [doi:10.1152/ajpregu.00628.2009](https://doi.org/10.1152/ajpregu.00628.2009)
- Kiyatkin, E.A., & Lenoir, M. (2011). Intravenous saline injection as an interoceptive signal in rats. *Journal of Psychopharmacology*. [doi:10.1007/s00213-011-2294-4](https://doi.org/10.1007/s00213-011-2294-4)
- Lenoir, M., & Kiyatkin, E.A. (2011). Critical role of peripheral actions of intravenous nicotine in mediating its central effects. *Neuropsychopharmacology*. Advance online publication. [doi:10.1038/npp.2011.104](https://doi.org/10.1038/npp.2011.104)
- Mazzone, G.L., & Nistri, A. (2011). Delayed neuroprotection by riluzole against excitotoxic damage evoked by kainate on rat organotypic spinal cord cultures. *Neuroscience*. Advance online publication. [doi:10.1016/j.neuroscience.2011.06.013](https://doi.org/10.1016/j.neuroscience.2011.06.013)
- Mazzone, G.L., & Nistri, A. (2011). Electrochemical detection of endogenous glutamate release from rat spinal cord organotypic slices as a real-time method to monitor excitotoxicity. *Journal of Neuroscience Methods*, 197(1): 128-132. [doi:10.1016/j.jneumeth.2011.01.033](https://doi.org/10.1016/j.jneumeth.2011.01.033)
- Muramatsu, T., Ohnuki, H., Ushio, H., Hibi, K., Igarashi, M., Hayashi, T., Ren, H., & Endo, H. (2011). Electrochemical flow injection immunoassay for cortisol using magnetic microbeads. *International Journal of Environmental and Analytical Chemistry*, 91(2): 161-173. [doi:10.1080/03067319.2010.500725](https://doi.org/10.1080/03067319.2010.500725)
- Smirnov, M.S., & Kiyatkin, E.A. (2010). Cocaine action on peripheral, non-monoamine neural substrates as a trigger of electroencephalographic desynchronization and electromyographic activation following i.v. administration in freely moving rats. *Neuroscience*, 165(2): 500-514. [doi:10.1016/j.neuroscience.2009.10.037](https://doi.org/10.1016/j.neuroscience.2009.10.037)
- Yasuzawa, M., Edagawa, K., Matsunaga, T., Takaoka, H., & Yabutani, T. (2011). Highly selective needle-type glucose sensors prepared by the immobilization of glucose oxidase on γ -polyglutamic acid film. *Analytical Sciences*, 27(3): 337-340. [doi:10.2116/analsci.27.337](https://doi.org/10.2116/analsci.27.337)